

Syddansk Universitet

**Data including GROMACS input files for atomistic molecular dynamics simulations of mixed, asymmetric bilayers including molecular topologies, equilibrated structures, and force field for lipids compatible with OPLS-AA parameters**

Róg, Tomasz; Orowski, Adam; Llorente, Alicia; Skotland, Tore; Sylvänne, Tuulia; Kauhanen, Dimple; Ekroos, Kim; Sandvig, Kirsten; Vattulainen, Ilpo

*Published in:*  
Data in Brief

*DOI:*  
[10.1016/j.dib.2016.03.067](https://doi.org/10.1016/j.dib.2016.03.067)

*Publication date:*  
2016

*Document version*  
Publisher's PDF, also known as Version of record

*Document license*  
CC BY-NC-ND

*Citation for pulished version (APA):*

Róg, T., Orowski, A., Llorente, A., Skotland, T., Sylvänne, T., Kauhanen, D., ... Vattulainen, I. (2016). Data including GROMACS input files for atomistic molecular dynamics simulations of mixed, asymmetric bilayers including molecular topologies, equilibrated structures, and force field for lipids compatible with OPLS-AA parameters. Data in Brief, 7, 1171-1174. DOI: 10.1016/j.dib.2016.03.067

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 09. Sep. 2018



ELSEVIER

Contents lists available at ScienceDirect

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](http://www.elsevier.com/locate/dib)

## Data Article

Data including GROMACS input files for atomistic molecular dynamics simulations of mixed, asymmetric bilayers including molecular topologies, equilibrated structures, and force field for lipids compatible with OPLS-AA parameters



Tomasz Róg<sup>a,b</sup>, Adam Orłowski<sup>a,c</sup>, Alicia Llorente<sup>d,e</sup>,  
Tore Skotland<sup>d,e</sup>, Tuulia Sylvänne<sup>f</sup>, Dimple Kauhanen<sup>f</sup>,  
Kim Ekroos<sup>f</sup>, Kirsten Sandvig<sup>d,e,g</sup>, Ilpo Vattulainen<sup>a,b,h,\*</sup>

<sup>a</sup> Department of Physics, Tampere University of Technology, P. O. Box 692, FI-33101 Tampere, Finland

<sup>b</sup> Department of Physics, University of Helsinki, P. O. Box 64, FI-00014, Finland

<sup>c</sup> Department of Physics and Energy, University of Limerick, Limerick, Ireland

<sup>d</sup> Department of Molecular Cell Biology, Institute for Cancer Research, Oslo University Hospital – The Norwegian Radium Hospital, 0379 Oslo, Norway

<sup>e</sup> Centre for Cancer Biomedicine, Faculty of Medicine, University of Oslo, 0379 Oslo, Norway

<sup>f</sup> Zora Biosciences Oy, 02150 Espoo, Finland

<sup>g</sup> Department of Biosciences, University of Oslo, 0316 Oslo, Norway

<sup>h</sup> MEMPHYS – Center for Biomembrane Physics, University of Southern Denmark, Odense, Denmark

## ARTICLE INFO

## Article history:

Received 8 December 2015

Received in revised form

6 March 2016

Accepted 18 March 2016

Available online 26 March 2016

## Keywords:

Molecular dynamics simulations

Force field

Topology

Lipidomics

Lipid

GROMACS

## ABSTRACT

In this Data in Brief article we provide a data package of GROMACS input files for atomistic molecular dynamics simulations of multi-component, asymmetric lipid bilayers using the OPLS-AA force field. These data include 14 model bilayers composed of 8 different lipid molecules. The lipids present in these models are: cholesterol (CHOL), 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphatidylcholine (POPC), 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphatidylethanolamine (POPE), 1-stearoyl-2-oleoyl-sn-glycero-3-phosphatidyl-ethanolamine (SOPE), 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphatidylserine (POPS), 1-stearoyl-2-oleoyl-sn-glycero-3-phosphatidylserine (SOPS), N-palmitoyl-D-erythro-sphingosyl-phosphatidylcholine (SM16), and N-lignoceroyl-D-erythro-sphingosyl-phosphatidylcholine (SM24). The bilayers'

DOI of original article: <http://dx.doi.org/10.1016/j.bbamem.2015.12.003>

\* Corresponding author at: Department of Physics, Tampere University of Technology, P. O. Box 692, FI-33101 Tampere, Finland.

E-mail address: [ilpo.vattulainen@tut.fi](mailto:ilpo.vattulainen@tut.fi) (I. Vattulainen).

<http://dx.doi.org/10.1016/j.dib.2016.03.067>

2352-3409/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

compositions are based on lipidomic studies of PC-3 prostate cancer cells and exosomes discussed in Llorente et al. (2013) [1], showing an increase in the section of long-tail lipid species (SOPS, SOPE, and SM24) in the exosomes. Former knowledge about lipid asymmetry in cell membranes was accounted for in the models, meaning that the model of the inner leaflet is composed of a mixture of PC, PS, PE, and cholesterol, while the extracellular leaflet is composed of SM, PC and cholesterol discussed in Van Meer et al. (2008) [2]. The provided data include lipids' topologies, equilibrated structures of asymmetric bilayers, all force field parameters, and input files with parameters describing simulation conditions (md.mdp). The data is associated with the research article "Interdigitation of Long-Chain Sphingomyelin Induces Coupling of Membrane Leaflets in a Cholesterol Dependent Manner" (Róg et al., 2016) [3].

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Specifications Table

Subject area	Chemistry, Biophysics
More specific subject area	Molecular dynamics simulations
Type of data	Text files
How data was acquired	Classical and quantum mechanics calculations
Data format	Consistent ordered package
Experimental factors	Software used: GROMACS 4 for MD, GAUSSIAN-03/09 for QM
Experimental features	OPLS-AA force field suit
Data source location	Not applicable
Data accessibility	Data are supplied with this article


Value of the data

- New parameters for MD simulations of PS and PE are provided.
- New parameters for MD simulations of sphingomyelin are provided.
- The lipid models given here are compatible with the OPLS-AA force field in studies of membrane proteins and drugs interacting with lipids.
- The force field parameters and lipid topologies provided here may be used to construct lipid bilayers whose composition is closely related to that of biological membranes.
- Equilibrated structures of asymmetric bilayer models may be used in MD studies of cell membranes.


1. Data

Lipidomics studies are becoming more and more popular [1] to identify in detail the lipid compositions found in cell membranes [2]. The emerging lipidomics data provide a very valuable source of information for atom-scale molecular dynamics (MD) simulations [3] that can clarify nanoscale dynamical phenomena in complex biomembrane systems.

To perform MD simulations, one needs detailed descriptions of the molecules that one aims to study [4]. First, one needs the topology, which includes information about all bonds, valence angles, and torsion angles in the molecule. Second, one needs a set of parameters, called the force field, to describe interactions related to all the interactions in the studied system. In this Data in Brief article, we provide a data package for the force field parameters and topologies for a number of lipids, in particular phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylserine (PS), and sphingomyelin (SM).



**Fig. 1.** Structure of SM16. The colors highlight the origin of force field parameterization for partial charges and torsional interactions. The color of the circle (red or blue) indicates the source of partial charges, and the color of the lines (red or black) connecting the circles indicates the source of torsional parameters. Red stands for parameterization based on [8], blue stands for parameterization based on new calculations reported in this work, and black corresponds to the original OPLS-AA parameters. There is reason to stress that all parameters for bonds and valence angle description are taken from the original OPLS-AA force field. Lennard-Jones parameters originate from OPLS-AA except for long hydrocarbons, which were derived in [8] (see text).



**Fig. 2.** Example of the simulation results: snapshots of equilibrated structures of a multicomponent bilayer (model M3 in Ref. [3]) where SM24 molecules are shown as yellow sticks with the last eight carbon atoms depicted as red balls. The figure illustrates interdigitation of long SM24 tails into the opposite leaflet of the bilayer. Lipids in the outer leaflet are shown as transparent blue glass, while lipids in the inner leaflet are described as transparent gray glass. For clarity, the system is surrounded by its periodic image on the left and right sides (SM24 molecules are highlighted only in the central image).

## 2. Experimental design, materials and methods

In this article, we provide the topologies and force field parameters for 6 lipids compatible with the OPLS-AA force field [5,6] to be used with the GROMACS simulation package [7]. The models described in this article are based on a combination of specific parameters for lipids derived in our previous studies [8–10], original OPLS-AA parameters, and a few specifically derived parameters for sphingomyelin. The description of cholesterol is entirely based on OPLS-AA parameters.

In our prior studies, we reparameterized (1) the torsion angles in the glycerol backbone and in the phosphocholine group, (2) the torsion angles and Lennard-Jones parameters in long saturated hydrocarbons [8], and (3) the torsion angles in unsaturated hydrocarbons [9]. All these components are necessary to describe a PC molecule. For PE and PS, we used the same parameters as for PC to describe the glycerol backbone, the phosphate group, and acyl chains, which are identical in these lipid classes. The parameters for the ethanolamine and serine groups were taken from the OPLS-AA force field derived for the amino acids' sidechain of lysine and serine, respectively.

The structure of SM is based on the amino alcohol sphingosine (Fig. 1) that is attached to a phosphatidylcholine head group and an acyl chain. To parameterize SM, we used previously derived parameters for the phosphatidylcholine group and long hydrocarbons [8], while the backbone part of sphingosine was reparameterized. First, partial charges were derived using a procedure described in our previous work [8], and the torsion angle potentials missing in the OPLS-AA set were calculated using a protocol described elsewhere [8]. Energy evaluation in torsion angles' potential parameterization was performed with the Hartree–Fock method by employing a polarized basis set 6–31G\* augmented with diffuse functions (6–31+G\*\*) on phosphorus and non-ester oxygen atoms. The effects of polar solvent were included implicitly by employing the Polarizable Continuum Model [11]. All electronic structure calculations were carried out using the GAUSSIAN-03/09 [12]. Fig. 1 shows the structure of SM16 with fragments of the molecule marked according to the source of parameters (Fig. 2).

## Acknowledgments

We thank Arkadiusz Maciejewski for performing QM calculations of rotational energy profiles and fitting the torsional potentials. We also thank CSC – IT Center for Science (Espoo, Finland) for computing resources. We further acknowledge the Academy of Finland (Center of Excellence program), the European Research Council (Advanced Grant CROWDED-PRO-LIPIDS), the Research Council of Norway through its Centers of Excellence funding scheme (Project no. 179571), the Norwegian Cancer Society, and the South-Eastern Norway Regional Health Authority for financial support.

## Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.03.067>.

## References

- [1] A. Llorente, T. Skotland, T. Sylvänne, T. Róg, D. Kauhanen, A. Orłowski, I. Vattulainen, K. Ekroos, K. Sandvig, Molecular lipidomics of exosomes released by PC-3 prostate cancer cells, *Biochim. Biophys. Acta* 1831 (2013) 1302–1309.
- [2] G. van Meer, D.R. Voelker, G.W. Feigenson, Membranes lipids: where they are and how they behave, *Nat. Rev. Mol. Cell Biol.* 9 (2008) 112–124.
- [3] T. Róg, A. Orłowski, A. Llorente, T. Skotland, T. Sylvänne, D. Kauhanen, K. Ekroos, K. Sandvig, I. Vattulainen, Interdigitation of long-chain sphingomyelin induces coupling of membrane leaflets in a cholesterol dependent manner, *Biochim. Biophys. Acta* 1858 (2016) 281–288.
- [4] I. Vattulainen, T. Róg, Lipid simulations: a perspective on lipids in action, *Cold Spring Harb. Perspect. Biol.* 3 (2011) a004655.
- [5] W.L. Jorgensen, D.S. Maxwell, J. TiradoRives, Development and testing of the OPLS all-atom force field on conformational energetics and properties of organic liquids, *J. Am. Chem. Soc.* 118 (1996) 11225–11236.
- [6] M.J. Robertson, J. Tirado-Rives, W.L. Jorgensen, Improved peptide and protein torsional energetics with the OPLS-AA force field, *J. Chem. Theory Comput.* 11 (2015) 3499–3509.
- [7] B. Hess, C. Kutzner, D. van der Spoel, E. Lindahl, GROMACS 4: algorithms for highly efficient, load-balanced, and scalable molecular simulation, *J. Chem. Theory Comput.* 4 (2008) 435–447.
- [8] A. Maciejewski, M. Pasenkiewicz-Gierula, O. Cramariuc, I. Vattulainen, T. Róg, Refined OPLS-AA force field for saturated phosphatidylcholine bilayers at full hydration, *J. Phys. Chem. B* 118 (2014) 4571–4581.
- [9] W. Kulig, M. Pasenkiewicz-Gierula, T. Róg, Cis and trans unsaturated phosphatidylcholine bilayers: a molecular dynamics simulation study, *Chem. Phys. Lipids* 195 (2016) 12–20.
- [10] W. Kulig, M. Pasenkiewicz-Gierula, T. Róg, Topologies, structures and parameter files for lipid simulations in GROMACS with the OPLS-aa force field: DPPC, POPC, DOPC, PEPC, and cholesterol, *Data Brief* 5 (2015) 333–336.
- [11] M. Cossi, G. Scalmani, N. Rega, V. Barone, New developments in the polarizable continuum model for quantum mechanical and classical calculations on molecules in solution, *J. Chem. Phys.* 117 (2002) 43–54.
- [12] M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, M.A. Robb, J.R. Cheeseman, J.A. Montgomery, T. Vreven, Jr., K.N. Kudin, J.C. Burant, et al., *Gaussian 03, Revision C.02*, Gaussian, Inc., Wallingford, CT, 2004.